



Salinity Stratification in the Upper Meter of the Ocean Observed During STRASSE


Bill Asher, Andy Jessup, Dan Clark
Applied Physics Laboratory
University of Washington
Seattle, Washington




Project funded by NASA Aquarius Program: Grant NNX09AU73G
With acknowledgment to Gilles Reverdin and the N/O *Thalassa*



The Surface Salinity Profiler



Towed Platform:
Measures S/T
outboard of ship
wake



4 x SBE-49
CTDs

5 cm →

20 cm →

50 cm →

100 cm →



100 cm




The Surface Salinity Profiler






Salinity Stratification in the Upper Ocean

STRASSE SSP Deployment 02




Low Wind



High $\Delta T_{\text{air-sea}}$
 Q_{lat} upwards


High Φ_{sw}






Time Series of Temperature and Conductivity From the SSP

Temperature





Conductivity






Upper Ocean Stratification Measured by the SSP





Density Stratification Measured by the SSP: Freshwater Flux: It is plausible?



From the bulk formula:

$$Q_{\text{Lat}} = \rho_A \Delta H_{\text{vap}} C_L U_{10} (q_W - q_A)$$


$$Q_{\text{Lat}} = 42 \text{ W/m}^2$$

From estimating mass loss:


$$Q_{\text{Lat}} = 39 \text{ W/m}^2$$



Salinity Stratification Measured by the SSP: Case 2, STRASSE SSP-10 09/10/2012




Low Wind




High $\Delta T_{\text{air-sea}}$
High Q_{Lat}

High Φ_{SW}






Salinity Stratification Measured by the SSP: Case 2, STRASSE SSP-10 09/10/2012







Salinity Stratification Measured by the SSP: Case 2

Temperature




Salinity








Salinity Contour at 50 cm from SSP data






Case 2: Temperature/Conductivity Details






Case 2: Salinity/Density Details

Salinity





Density





Case 2: Salinity Gradients



Well-mixed salinity vs. stratified salinity region





Case 2: Density Gradients

Well-mixed salinity vs. stratified salinity region





STRASSE Data Analysis: Further Work

Implement 1-D ocean mixed layer model to incorporate mixing, verify that positive salinity gradients can form due to evaporation with high insolation

Continue analysis of freshwater gradients measured in E. Equatorial Pacific